How does COVID-19 affect the heart? New clues arise



Written by <u>Annie Lennon</u> on August 1, 2022 — <u>Fact checked</u> by Alexandra Sanfins, Ph.D.



How, exactly, doees SARS-CoV-2 cause damage to the heart? Image credit: Stefania Pelfini, La Waziya Photography/Getty Images.

- Researchers investigated how SARS-CoV-2 damages heart muscle cells.
- They found that the virus damages heart muscle cells via an inflammatory pathway independent of ACE2 receptors.
- They now plan to investigate how SARS-CoV-2 causes inflammation in the heart.

Studies show that up to $62\%^{\circ}$ of patients hospitalized with COVID-19 experience cardiac injury.

Other <u>research</u> shows that individuals who have contracted COVID-19-

Knowing more about how SARS-CoV-2 affects the heart could help develop new treatments to protect cardiac health.

Recently, researchers conducted a series of experiments that suggested SARS-CoV-2's spike protein can lead to heart muscle injury through an inflammatory process.

"It's already known from the clinical side that COVID-19 infection can induce heart injury, however, what we don't know is the mechanistic details of how this occurs. What we suspect is that the spike protein has unknown pathological roles," said <u>Dr. Zhiqiang Lin</u>, lead author of the study and an assistant professor at the Masonic Medical Research Institute in Utica, New York.

"Our data show that the spike protein from SARS-CoV-2 causes heart muscle damage. That's why it's important to get vaccinated and prevent this disease," he added.

The researchers presented their findings at the American Heart Association's <u>Basic Cardiovascular Sciences Scientific Sessions 2022</u>[©].

ADVERTISEMENT

Q

MEDICALNEWSTODAY

SUBSCRIBE

The role of the spike protein

SARS-CoV-2 enters healthy cells via its spike protein, located on its surface. The spike protein works by latching onto receptors known as

<u>HCoV-NL63</u> is also a coronavirus that enters cells by attaching to ACE2 receptors. While it causes moderate respiratory symptoms, it does not lead to cardiac injury.

"Host natural immunity is the first line of defense against pathogen invasion, and heart muscle cells have their own natural immune machinery. Activation of the body's immune response is essential for fighting against virus infection; however, this may also impair heart muscle cell function and even lead to cell death and <u>heart failure</u>," explained Dr. Lin.

In the present study, the researchers hypothesized that SARS-CoV-2 may damage the heart via immune responses that are independent of ACE2. To test their hypothesis, they compared the ability of both viruses to cause cardiac injury.

They found that while SARS-CoV-2 activated the <u>TLR4 signaling pathway</u> • – a major part of the natural immune response — while HCov-NL63 did not.

To understand more about how activation of TLR4 signaling may affect the heart, the researchers cloned the SARS-CoV-2 spike protein and the NL63 spike protein, and delivered them to mice via an <u>adeno-associated</u> <u>viral vector</u>.

They found that the SARS-CoV-2 spike protein, and not the NL63 protein, led to heart dysfunction, hypertrophic remodeling— or in other words, cellular enlargement—and cardiac inflammation.

In vitro experiments with rat cardiac muscle cells also led to hypertrophic remodeling and increased levels of inflammatory markers.

Heart damage from the spike protein

The researchers next examined heart biopsies from a deceased patient with COVID-19-associated myocarditis, a patient who died of mRNA-

Whereas both the spike protein and the TLR4 protein were detected in cardiac muscle cells and other cell types in the COVID-19 patient, neither protein was detected in the healthy heart.

"That means once the heart is infected with SARS-CoV-2, it will activate TLR4 signaling. Besides directly damaging the heart muscle cells, the spike protein itself is very inflammatory and may cause systemic inflammation that indirectly causes heart problems," noted Dr. Lin.

Meanwhile, the researchers detected only a small fraction of spike protein in the cardiac muscle cells of the mRNA vaccine patient and no obvious signs of TLR4.

This, they noted, means that mRNA vaccines are unlikely to cause myocarditis via cardiac muscle cells.

The researchers concluded that the SARS-CoV-2 spike protein might damage the heart via TLR4 inflammatory pathways, independently of ACE2.

ADVERTISEMENT

Study limitations

When asked about possible limitations to the study, <u>Dr. Ziyad Al-Aly</u>, chief of research and development at VA Saint Louis Health Care System, told *MNT*: "This is preliminary data in the form of an abstract, so it is hard to

inquiry is important to help us understand how SARS-CoV-2 affects the heart, and what role, if any, spike proteins may play."

"It also helps us understand what the mechanisms of injury may be. Understanding these mechanisms will help design preventive and therapeutic strategies," he added.

<u>Dr. James Lo</u>, assistant professor at Weill Cornell Medicine Graduate School of Medical Sciences, agreed that it is difficult to comment on the study's limitations without seeing the full paper.

He nevertheless called the findings "intriguing," and noted:

"What is still unknown is how the spike protein would activate inflammatory pathways such as TLR4. We still have a long way to go in terms of understanding why a small minority of patients with COVID-19 develop cardiac injury or myocarditis."

Moving forward, the researchers intend to investigate how SARS-CoV-2 spike proteins cause inflammation in the heart.

So far, they have outlined two potential ways this may happen. The first possibility is that the spike protein directly activates inflammation in virus-infected heart muscle cells.

The second possibility is that the spike protein is shed into the bloodstream and damages the heart as it circulates.

ADVERTISEMENT

Check your vitamin levels with a micronutrient test

This micronutrient test checks for vitamin B12, D, and E, plus magnesium,

conner selenium and zinc Get vour results in 2-5 days from an

Heart Disease Infectious Diseases / Bacteria / Viruses COVID-19



FEEDBACK:

RELATED COVERAGE

COVID-19 and the heart: What do we know so far?

SARS-CoV-2 affects various parts of the body, including the cardiovascular system. In this feature, we examine the latest research an...

READ MORE

In Conversation: Long COVID's cardiovascular implications

Three experts and one long hauler join us in conversation to explain long COVID's cardiovascular impact and postural orthostatic tachycardia...

READ MORE

Why does the coronavirus cause blood

This article will discuss the blood clotting process, analyze the latest research on why the coronavirus causes blood clots, and offer some tips...

READ MORE

Is chest pain due to COVID-19, anxiety, a heart attack, or another cause?

Medically reviewed by Cameron White, MD, MPH

Chest pain can sometimes be a sign of COVID-19, anxiety, or a heart attack. Learn about the treatment options for each of these causes and...

READ MORE

f ¥

Get our newsletter

Keep up with the ever-changing world of medical science with new and emerging developments in health.

Enter your email

SUBSCRIBE

Your privacy is important to us

ADVERTISEMENT

Contact Us	Health Hubs
Terms of Use	Medical Affairs
Privacy Policy	Content Integrity
Privacy Settings	Newsletters
Advertising Policy	Do Not Sell My Info

© 2004-2022 Healthline Media UK Ltd, Brighton, UK, a Red Ventures Company. All rights reserved. MNT is the registered trade mark of Healthline Media. Any medical information published on this website is not intended as a substitute for informed medical advice and you should not take any action before consulting with a healthcare professional. See additional information.

About | Careers | Advertise with us

OUR BRANDS

Medical News Today Greatist Psych Central